Quality Harvest



nutrimaster

Grow like a Pro

Cabbage



Mike +263 718 787 677 , **Tiaan** +263 711 900 765
Josh +263 788 270 997**, Loyd** +263 0719 124 673
Visit us at 17 Foundry Road, Aspindale, Harare.

Scan me

Optichem
OPTIMISE YOUR GROWTH

Cabbage Programme

Products	Control	Days After Planting	Rate/ha or per 100lts Water	Comments
Nutrition				
Cereal Blend 6.23.23	Nutrition	0 days	400kgs	Apply as a band into side of ridge before transplanting take care not to plant seedling directly into fertilizer band this is a general recommendation always follow soil sample results
Ammonium Nitrate	Nutrition	Day 7, Day 21, Day 35	50 kg/ha per application (150 kg/ha total)	Apply as split applications to supply nitrogen during early vegetative growth.
Nutristarter	Nutrition	7	4lt	Apply as a foliar once active growth is seen in plant only one application
K Silica	Nutrition	Day 7, Day 14, Day 21, Day 28, Day 35	2 L/ha per application (10 L/ha total)	Apply as a foliar once active growth is seen; repeat weekly on the dates shown.
APP	Nutrition	7 and 21	15lt	Apply as a foliar once active growth is seen in plant apply second application 2 weeks after 1st
Nutri Grow	Nutrition	Day 14 and Day 21	4lt	Apply as a foliar at Week 2 and Week 3 after planting.
Nutri Bloom	Nutrition	Day 35 and Day 42	4lt	Support head initiation and early head development.
Foliariel OK	Nutrition	35	500g	Apply at the start of head initiation to improve head firmness and reduce calcium deficiency.
Calcium nitrate	Nutrition	Day 35 and Day 49	87.5 kg/ha per application (175 kg/ha total)	Apply at the start of head initiation to improve head firmness and reduce calcium deficiency.
Sop	Nutrition	Day 49	250kg	Apply at the start of head formation to support potassium demand and head development.
Herbicides				
Glyphosate Xtra 41%	Pre emergence Non selective herbicide	14 days pre planting	4lts	Pre irrigate so as to encourage weed growth 28 days before planting Apply glyphosate at 14 days before planting. Ensure ridges are up before this so that herbicide is not distrubed by cultivation of soil.
Fungicides				
Macometa	Downy Mildew.	7 and 21	280g/100lts	Maximum of 4 Applications per season
Copper Hydroxide	Downy Mildew.	14 and 28	100mls/100lts	Maximum of 6 applications per season
Insecticides				
Worm kill	Caterpillars	First sign of pest	105-150ml	Maximum of 2 applications per season
Emmamectin duo	Caterpillars	First sign of pest	75g/100lts	Apply at first sign of pest
Cartap	Caterpillars	First sign of pest	400g/100lts	Apply at first sign of pest maximum 2kg/ha
				Apply at first sign of pest
				Apply at first sign of pest
				Apply at first sign of pest. Do not exceed 3 applications per season

PLEASE READ CHEMICAL LABELS BEFORE APPLICATION



