Quality Harvest



nutrimaster

Grow like a Pro

Peas



Mike +263 718 787 677 , **Tiaan** +263 711 900 765
Josh +263 788 270 997, **Ethan** +263 783 835 724
Visit us at 17 Foundry Road, Aspindale, Harare.

Scan me

Opti**chem**
OPTIMISE YOUR GROWTH

Peas Programme

Products	Control	Days After Planting	Rate/ha or per 100lts Water	Comments
Nutrition				
K Silica	Cell wall strengthening	14days, 28days, 42days	2	
APP - Ammonium Poly Phosphate	Nutrition	at planting - 3 applications 2 weeks apart	15	
Nutristarter	Vegatative Nutrition	14-21	4lt	Apply as a full cover spray every 7 – 14 days just after germination or establishment of transplants.
Nutri Grow	Vegatative Nutrition	28-35	4lt	Apply as a full cover spray at 1 – 2-week intervals to promote vegetative growth. Application range of 1-4L can be used.
Nutri Bloom	Flowering	42-49	4lt	Nutribloom to assist with bulking period or during periods that require high Nitrogen levels.
Trichotech	Seed dressing	0	125g/ha of seed	Fusarium, Pythium, Rhizoctonia control.
Mytech	Seed dressing	0	125g/ha of seed	
Foliariel OK	Nutrition	At planting	200g/ha	Apply as a drench or with Knapsack to the surface of the bed prior to planting.
Zinc Chelate	Nutrition	At planting	300g/ha	Apply as a drench or with Knapsack to the surface of the bed prior to planting.
Copper Chelate	Nutrition	At planting	300g/ha	Apply as a drench or with Knapsack to the surface of the bed prior to planting.
Compound 6.23.23	Nutrition	0	500kg/ha	
Ammonium Nitrate	Nutrition	20-50	250kg/ha	
MOP	Nutrition	49-59	150kg/ha	
Calcium Nitrate	Nutrition	6	50kg/ha	
Herbicides				
Glyphosate Xtra 41%	Pre-emergence	14 days pre planting	2lts	Pre irrigate so as to encourage weed growth 28 days before planting Apply glyphosate at 14 days before planting. Ensure ridges are up before this so that herbicide is not distrubed by cultivation of soil.
Fusilade	Post-emergence grass control	14 days post planting	2.5lts	
Bentazone 480SL	Post-emergence broadleaf control	14 days post planting	3lts	
Fungicides				
Amistar top	Ascochyta, Powdery mildew	21 days and 7 day intervals after first application	500ml	Apply as a preventative spray do not apply more than 6 applications per season.
Folicur	Powdery mildew	21 days	50ml/100lts water	500l spray mix per ha at first sign of disease and then 10-14 day intervals after.
Bravo	Ascochyta	35-45 days short season variety 50-60 days long season varieties.	1.4lts	Maximum of 3 sprays per season.
Insecticides				
Nemshield	Nematodes	7 days pre planting	2 lts a hectare	Maximum of 2 applications per season.
Coragen	Heliothis Bollworm (larvae)		100 mL/ha for bollworm control	Apply at first sign of pest.
Karate Zeon	Boring pests	At first sign of pest	200ml	In 200 to 1000 lts of water 1-4 sprays
Beauvitech	Thrips and white fly		20g/100lts	
Lecatech	Whitefly		250g	
Acela 60ec	Leafminer, aphids, thrips		90ml/100lts	

PLEASE READ CHEMICAL LABELS BEFORE APPLICATION