

nutrimaster

Grow like a Pro

Tomatoes



Mike +263 718 787 677 , **Tiaan** +263 711 900 765
Josh +263 788 270 997, **Ethan** +263 783 835 724
Visit us at 17 Foundry Road, Aspindale, Harare.

Scan me

Optichem
OPTIMISE YOUR GROWTH

Tomato Programme Proposal

Products	Control	Days After Planting	Rate/ha or per 100lts Water	Comments
Nutrition				
Compound C 6.23.23	Nutrition	0	400kg	Apply as a band into side of ridge before transplanting take care not to plant seedling directly into fertilizer band this is a general recommendation always follow soil sample results.
Ammonium Nitrate	Nutrition	7 to 35	150kg	Broadcast over bed once active plant growth is seen split applications of 50kg per week from day 7 after planting.
Nutristarter	Nutrition	7	4lt	Apply as a foliar once active growth is seen in plant only one application.
K Silica	Nutrition	7 to 35	2lt	Apply as a foliar once active growth is seen in plant repeat weekly.
APP	Nutrition	7 to 21	1.5lt	Apply as a foliar once active growth is seen in plant apply second application 2 weeks after 1st.
Nutri Grow	Nutrition	21 and 28	4lt	Apply as a foliar at week 2 after planting and week 3.
Nutri Bloom	Nutrition	35 and 42	4lt	To assist in the flowering process.
Foliarel OK	Nutrition	35	500g	Mix with first application of Nutri Bloom.
Calcium Nitrate	Nutrition	35 and 39	175kg	Apply 100 kg/ha at frist fruit set follow up with 75kg 3 weeks later.
SOP	Nutrition	49	200kg	Apply as a soil application from start of fruit set to the end of fruit set splitting application weekly
Herbicides				
Glyphosate Xtra 41%	Pre emergence Non selective herbicide	14 days before planting	4lts	Cultivation of Soil.
Metribuzin 480 SC	Post emergence Broadleaf control	14	1.5lts	Apply to actively growing weeds.
Fusilade	Post Emergence Grass Control	14	2.5lts	Apply to actively growing weeds.
Fungicides				
Macometa	Late Blight control	7 and 21	280g	Maximum of 4 Applications per season.
Opti stroby	Early Blight Control	14 and 28	500ml	Maximum of 6 Applications per season.
Copper Oxychloride	Bacterial spot	7 to 49	500g	Weekly applications from 7 days after planting when crop is actively growing.
Insecticides				
Imidacloprid	White fly and Aphids	7	30mls/100lts	Maximum of 2 applications per season.
Acetamiprid	White fly, American leaf miner and Aphids	14	50g/100lts	Apply at first sign of pest.
Coragen	White fly, American leaf miner and Aphids	14	20mls/100lts	Apply at first sign of pest.
Optimator	White fly, American leaf miner, Aphids and Thrips	14	100mls/100lts	Apply at first sign of pest.
Tuta Guard	Fruit Borer	49	20mls/100lts	Apply at first sign of pest.
Emmamectin duo	Boll worm and American Leaf miner	49	150g	Apply at first sign of pest. Do not exceed 3 applications per season.

PLEASE READ CHEMICAL LABELS BEFORE APPLICATION